GENERAL

# management of a pediatric unit hesi case study

Published: September 3, 2026 | Index ID: #-

High Expectation Student Assessment (HESI) exams are a cornerstone of nursing education, ensuring that new graduates possess the knowledge and skills required for safe patient care. For those entering the pediatric field, the HESI Pediatric Nursing exam adds an extra layer of complexity, testing familiarity with child specific physiology, pharmacology, and developmental considerations. This article presents a detailed case study that illustrates the **management of a pediatric unit HESI case study**, offering a step by step roadmap for educators, clinical preceptors, and students preparing for the exam.

## Why Focus on a Pediatric Unit HESI Case Study?

In the fast paced environment of a pediatric unit, nurses must juggle acute medical conditions, developmental milestones, and family dynamics. The HESI exam reflects this reality by probing not only factual recall but also critical thinking and application in clinical scenarios. By dissecting a realistic case study, learners can:

- Understand the structure of HESI questions and the typical themes tested.
- Translate textbook knowledge into bedside practice.
- Identify gaps in their preparation and target study efforts.
- Develop strategies for time management and stress reduction during the exam.

## Structure of the Case Study

Our case study follows a standard HESI format: a patient vignette, a series of multiple choice questions, and a set of short answer prompts. The case is designed to mimic a real pediatric admission, incorporating common conditions such as asthma exacerbation, viral bronchiolitis, and pediatric sepsis.

## Case Overview

### Patient Profile

**Name:** Emma Martinez**Age:** 7 years old**Chief Complaint:** Shortness of breath, wheezing, and cough for 3 days **History:** Known atopic asthma, well controlled on albuterol inhaler, no recent hospitalization**Family History:** Mother has allergic rhinitis, father has hypertension**Allergies:** None documented

### Clinical Presentation

Emma is visibly distressed, clutching her chest. She is using accessory muscles, has a rapid respiratory rate of 32 breaths per minute, and oxygen saturation is 88% on room air. Auscultation reveals bilateral wheezes and decreased air entry. She is afebrile, heart rate 110 bpm, blood pressure 95/60 mmHg.

### Initial Assessment Findings

- Peak expiratory flow (PEF) measured at 350 L/min (50% of predicted)
- Peak inspiratory flow (PIF) 70 L/min
- Chest X ray pending, but preliminary imaging suggests hyperinflation.

### Laboratory Results

1. Complete blood count (CBC): White blood cell count 12,000/µL, eosinophils 6%

2. Arterial blood gas (ABG): pH 7.35, PaCO , 45 mmHg, PaO , 65 mmHg, HCO ƒ { 22 mEq/L
3. Serum electrolytes: Na z 140 mEq/L, K z 4.2 mEq/L, Cl { 104 mEq/L, HCO ƒ { 24 mEq/L
4. Blood glucose 95 mg/dL

## Key HESI Themes Covered in the Case

- Pharmacology: Appropriate drug selection, dosing, and monitoring for pediatric asthma.
- Pathophysiology: Understanding airway inflammation, bronchoconstriction, and the role of eosinophils.
- Assessment & Monitoring: Interpretation of peak flow readings, ABG, and vital signs.
- Patient & Family Education: Communicating asthma action plans and inhaler technique.
- Critical Thinking: Recognizing when to elevate care and anticipate complications.

## Step by Step Management of the Pediatric Unit HESI Case Study

### 1. Pre Assessment Preparation

Before reviewing the case, students should:

- Review core pediatric concepts: growth charts, developmental milestones, and common illnesses.
- Study the pharmacology of bronchodilators, steroids, and antileukotrienes.
- Practice interpreting peak flow ratios and ABG values.
- Familiarize themselves with HESI question patterns: “most appropriate next step,” “best explanation,” and “risk factors.”

### 2. Detailed Patient Assessment

Using the **APGAR assessment for pediatrics**(Airway, Breathing, Circulation, Assessment, and Response), evaluate Emma’s status:

- Airway: Patent, no obstruction.
- Breathing: Rapid, using accessory muscles.
- Circulation: Tachycardic, borderline hypotension.
- Assessment: Decreased oxygen saturation.
- Response: Distressed, crying.

### 3. Prioritize Interventions

1. Administer a short acting beta agonist (SABA) via nebulizer: 2.5 mg albuterol.
2. Begin supplemental oxygen to maintain SpO , > 94%.
3. Reassess peak flow and ABG after 20 minutes.
4. If no improvement, consider systemic corticosteroids (e.g., oral prednisone 0.5 mg/kg).
5. Monitor for signs of impending respiratory failure: worsening tachypnea, decreased consciousness.

### 4. Documentation and Communication

Accurate, timely documentation is critical. Use the **SBAR (Situation, Background, Assessment, Recommendation)** format for handoffs:

- Situation: Emma Martinez, 7 year old, acute asthma exacerbation.
- Background: History of atopic asthma, well controlled.
- Assessment: SpO , 88%, PEF 350 L/min.
- Recommendation: Initiate nebulized albuterol, supplemental O ,, and monitor response.

### 5. Family Engagement

Effective communication with Emma’s parents is essential for compliance and education:

1. Explain the severity of the exacerbation and the need for aggressive treatment.
2. Review inhaler technique and demonstrate correct use.
3. Distribute an updated asthma action plan.
4. Encourage parents to monitor peak flow at home and seek emergency care if readings drop below 70% predicted.

### 6. Follow Up and Discharge Planning

Once Emma stabilizes (SpO , > 94%, PEF > 70% of predicted, and no accessory muscle use), plan for discharge:

- Provide a written asthma action plan with medication schedules.
- Schedule a follow up visit with the pediatric pulmonologist in 2 weeks.
- Arrange for home health nurse evaluation if needed.
- Educate on avoidance of known triggers (e.g., dust mites, pet dander).

## Sample HESI Style Questions Derived from the Case

### Multiple Choice Question 1

**What is the most appropriate next step in Emma's management?**

1. Administer a short acting beta agonist via nebulizer.
2. Begin oral prednisone immediately.
3. Order a chest X ray.
4. Provide supplemental oxygen via nasal cannula at 2 L/min.

**Correct Answer:** A. Administer a short acting beta agonist via nebulizer.

### Short Answer Prompt

**Explain why a peak flow ratio < 70% of predicted is significant in pediatric asthma management.**

**Answer:**A peak flow ratio below 70% of predicted indicates severe airway obstruction, correlating with increased risk of exacerbation and the need for intensified therapy. It guides clinicians to adjust medication regimens and monitor for potential deterioration.

## Common Pitfalls in Pediatric HESI Preparation

- Overemphasis on pharmacology at the expense of assessment skills.
- Neglecting the developmental context when interpreting symptoms.
- Failing to practice time management strategies for the timed exam.
- Ignoring the importance of patient and family education.
- Underestimating the role of interprofessional collaboration.

## Strategies to Overcome These Pitfalls

### 1. Integrate Clinical Scenarios into Study Sessions

Use case based learning modules that require students to apply knowledge in realistic contexts, mirroring the HESI exam structure.

### 2. Focus on Developmental Milestones

Incorporate developmental checklists into your study routine, ensuring you can differentiate between age appropriate behaviors and signs of pathology.

### 3. Time  Management Drill Runs

Simulate exam conditions with timed quizzes, gradually decreasing the allowed time per question to build speed and confidence.

### 4. Role  Playing Family Education

Practice explaining complex medical information to a “family member” (peer or tutor) to improve communication skills and reduce anxiety during actual patient interactions.

### 5. Build a Collaborative Study Group

Regularly meet with peers to discuss challenging questions, share resources, and provide mutual support.

## Interprofessional Collaboration in the Pediatric Unit

Effective pediatric care relies on a multidisciplinary team. Key collaborators include:

- Pediatricians: Provide overall medical oversight.
- Respiratory Therapists: Optimize inhalation therapy and monitor ventilatory status.
- Pharmacists: Ensure appropriate dosing and drug interactions.
- Social Workers: Address socioeconomic barriers to care.
- Physical & Occupational Therapists: Assist with mobility and developmental support.

In the HESI case, the nurse must coordinate with the respiratory therapist for nebulizer setup and with the pharmacist for accurate dosing calculations.

## Documentation Standards: The Cornerstone of Quality Care

High  quality documentation serves several purposes:

- Provides a legal record of care.
- Facilitates continuity during handoffs.
- Supports data collection for quality improvement.
- Demonstrates compliance with regulatory standards.

Use the **SOAP (Subjective, Objective, Assessment, Plan)** format for progress notes:

1. Subjective: Emma reports “I feel like I can’t breathe.”
2. Objective: SpO , 88%, PEF 350 L/min.
3. Assessment: Acute asthma exacerbation.
4. Plan: Nebulized albuterol, O ,, monitor response.

## Challenges and Solutions in Pediatric Unit Management

### Challenge 1: Rapid Patient Deterioration

Children can decompensate quickly, necessitating immediate escalation.

**Solution:** Maintain a low threshold for calling

## Baca Juga (Artikel Terkait)

- **automaker mini software version v8 erptoboss**
  URL: http://www.directnilecruiseegypt.com/automaker-mini-software-version-v8-erptoboss.pdf
- **automata and computability**
  URL: http://www.directnilecruiseegypt.com/automata-and-computability.pdf
- **automata and languages theory and applications**
  URL: http://www.directnilecruiseegypt.com/automata-and-languages-theory-and-applications.pdf
- **automata and mechanical toys**
  URL: http://www.directnilecruiseegypt.com/automata-and-mechanical-toys.pdf

**Tags: #management #pediatric #unit #hesi #case #study**